

**The Swedish Sea Level Network**  
GLOSS Experts 12<sup>th</sup> Meeting, November 2011  
Thomas Hammarklint

Swedish Meteorological and Hydrological Institute,  
Sven Källfelts gata 15, SE-42671 Göteborg, Sweden  
Telephone: +46 11 4958000, Email: [Thomas.Hammarklint@smhi.se](mailto:Thomas.Hammarklint@smhi.se)

## Introduction

The Swedish Sea Level Network, operated by the Swedish Meteorological and Hydrological Institute (SMHI), records sea level at 23 locations (Figure 1 and Table 1). The network is considered as the official Swedish sea level network. SMHI is responsible both for the network, data and the levelling of the stations. The Swedish sea level records constitute some of the longest and most robust sea level records in the world. Also, the Swedish Maritime Administration, records sea level at about 30 locations. A list of these stations is shown in Appendix 1.




**Figure 1.** The Swedish Sea Level Network operated by SMHI, November 2011.



**Figure 2.** Two Swedish GLOSS-stations; Göteborg-Torshamnen and Stockholm-Skeppsholmen.

## The Swedish sea level network

The first systematic Swedish observations of the sea level started 1774 at the sluice in the harbour of Stockholm. At the end of 19<sup>th</sup> century the Swedish king decided to establish seven mareographs, where several are still operating or have been substituted by other stations. In 1889 the Nautical-Meteorological Bureau (a predecessor of SMHI) established a continuously recording sea level station in the bedrock (mareograph) on the island Skeppsholmen, located close to the sluice. This mareograph has since then recorded the Stockholm sea level and is now operated by SMHI. The sea level series in Stockholm constitutes the longest sea level record in the world (Figure 3).




**Figure 3.** Annual mean sea levels in Stockholm since 1774, with the regression line (corresponding to land uplift) for 1774-1885 and its extension into modern times. The increased sea level rise since the late 19<sup>th</sup> century appears as a deviation from the regression line.

During the 20<sup>th</sup> century more stations were established. The technique used from the beginning was the stilling well technique. The Sea Level Network was completely modernised during the 1980s. The traditional stilling well was still used, but the gauges were converted from analogue to digital with automatic data transfer to SMHI. Earlier the recording was only done with a chart recording apparatus. This mechanical equipment is nowadays used as a backup for the digital recording equipment, mainly to prevent gaps in the time-series.

A new modernisation of the network was completed at the end of 2005. A new data logger (Vaisala MAWS) was installed that is more capable of delivering near real time data. The data recorded by the measurement equipment is transferred to SMHI once an hour through the telephone line and stored in a database. From there, the data can be presented in real-time on our website and in our FTP-box. Quality controlled data are distributed to users via national and international exchange on a continuous basis. The software MATLAB is used for validation and correction of the data. We can fill gaps with data from paper charts or predictions and add or subtract a constant offset to the data. The original data are stored in a separate table in the database.

At present we store data 10-minute values and also the maximum and minimum records every hour are stored. In the future we will store all data recorded by the equipment (one minute values) in our database. In order to check the status of a station and validate real-time data an observer visits the station once a week. The sea level station is connected to several Bench Marks. The Swedish mapping, cadastral and land registration authority (Lantmäteriet) does the precise levelling, i.e. they are responsible for determining the distance between the Contact Point and the Bench Marks. SMHI is responsible for keeping Tide Gauge Zero (TGZ) a fixed distance below the Contact Point. Most of the gauges are installed in the bedrock, but some are located in slightly unstable areas. Levelling is done once a year. The levelling often shows no significant vertical motion on the majority of the sea level stations.

Figure 4 shows the basic structure of a typical sea level station (mareograph). Sea level is measured in a deep well beneath the mareograph building. The well is connected to the sea through a narrow underwater pipe, to damp out short-period fluctuations of the sea level.



**Figure 4.** Basic structure of a typical sea level station or mareograph.

The mechanical part of the measurement equipment is constructed of a float, floating on the water surface, connected to a counterweight with a steel band. The steel band is attached on a wheel, which is connected to the digital equipment. When the sea level varies and the float follows it up and down, the equipment registers the rotation of the wheel, which is transformed into a digital reading using an encoder (Vaisala QSE 102).

Station	Latitude	Longitude	Digital data available from	Installation and type of CGPS	Distance CGPS (km)	Installation of AG
Kalix	65° 41' 49" N	23° 05' 46" E	1974	No	-	No
Furuögrund	64° 54' 57" N	21° 13' 50" E	1916	1993A	9.5	1992
Ratan	63° 59' 10" N	20° 53' 42" E	1891	2006A	0.1	2007
Skagsudde*	63° 11' 26" N	19° 00' 45" E	1982	No	-	No
Spikarna	62° 21' 48" N	17° 31' 52" E	1968	No	-	No
Forsmark	60° 24' 31" N	18° 12' 39" E	1975	No	-	No
Stockholm-Skeppsholmen	59° 19' 27" N	18° 04' 54" E	1889	1992A/B	15.3/3.4	No
Landsort Norra	58° 46' 08" N	17° 51' 32" E	2004	No	-	No
Marviken	58° 33' 13" N	16° 50' 14" E	1964	No	-	No
Visby*	57° 38' 21" N	18° 17' 04" E	1916	1993A	5.2	2004
Ölands norra udde*	57° 21' 58" N	17° 05' 50" E	1851	2004B	13.5	No
Oskarshamn	57° 16' 30" N	16° 28' 41" E	1960	No	-	No
Kungsholmsfort	56° 06' 19" N	15° 35' 22" E	1886	2004A	0.1	No
Simrishamn	55° 33' 27" N	14° 21' 28" E	1982	No	-	No
Skånör	55° 25' 00" N	12° 49' 46" E	1992	2002B	1.8	No
Klagshamn	55° 31' 20" N	12° 53' 37" E	1929	No	-	No
Barsebäck	55° 45' 23" N	12° 54' 12" E	1937	2002B	5.9	No
Viken	56° 08' 32" N	12° 34' 45" E	1976	No	-	No
Ringhals	57° 14' 59" N	12° 06' 45" E	1967	1991A	19.7	1993
Göteborg-Torshamnen	57° 41' 05" N	11° 47' 26" E	1967	2004B	12.8	1976
Stenungsund*	58° 05' 36" N	11° 49' 57" E	1962	No	-	No
Smögen	58° 21' 13" N	11° 13' 04" E	1910	2002A	0.05	2004
Kungsvik	58° 59' 48" N	11° 07' 38" E	1976	2005B	7.4	No

**Table 1.** List of stations in the Sea Level Network operated by SMHI. Stations marked \* are non-realtime reporting stations. CGPS marks places where Continuous Global Positioning is installed and measurements of the absolute land uplift are being carried out. Type of CGPS: A denotes complete stations (EUREF reference stations with antennas placed on solid bedrock), B simplified stations (mounted on buildings). Distance CGPS means the distance between the CGPS antenna and the sea level station. Only CGPS-stations located less than 20 km from a sea level station are included. AG means that the station has a platform for observing Absolute Gravity. More sealevel data is available from discontinued stations. Some stations were located near the existing stations.

## Co-location of geodetic observing system at mareographs

Lantmäteriet has developed the geodetic infrastructure at several of the mareographs to include connection to the national height levelling network, continuous GNSS as well as absolute gravity. The main purpose of these techniques has been to develop a model to describe the post glacial rebound. One of the main tasks for the geodetic research division at Lantmäteriet is to develop, monitor and maintain the national reference systems and frames in all dimensions (3D, horizontal, height) as well as gravity so that the need of the society is satisfied.

The national levelling network was levelled during the third precise levelling of Sweden during 1978-2001 and resulted in the height system RH2000, which is the Swedish realization of the European height system EVRS. GNSS at mareographs was first done as a GPS-campaign during the European project EUVN in 1997. The monuments have later been equipped with CGPS, see Table 1, and are now part of the Swedish CORS network named SWEPOS™.

Lately, several different Nordic institutions as well as other international actors have observed gravity with absolute gravimeters in the Nordic and Baltic area. These efforts have been co-ordinated through the working group of geodynamic within NKG (Nordic Commission of Geodesy). The main purpose of these measurements has been to detect the change of gravity over time, mainly caused by the post glacial rebound. Several mareographs are today equipped with an absolute gravity platform (Figure 5).



**Figure 5.** Smögen, a mareograph (hut to the left) also combined with CGPS (monument to the right) and absolute gravity platform (hut in the middle).


## Historical sea level data

In November 2011 the sea level database at SMHI contained more than 3000 years with digital sea level observations, where about 1600 years are from continued stations. Most of the data are hourly values, but for the past years, the resolution has been increased to 10-minute values. A complete station list showing the content of the data base on a yearly basis can be found here:

[http://www.smhi.se/hfa\\_coord/BOOS/dbkust/Availability\\_Sealevel\\_SMHI.htm](http://www.smhi.se/hfa_coord/BOOS/dbkust/Availability_Sealevel_SMHI.htm)

## Climate changes in sea level data

From our long time series we can detect the global sea level rise after reducing the yearly means with the land-uplift effect (Figure 6). A regression analysis indicates a sea level rise around 3 mm per year for the last 30 years and approximately 1.5 mm per year since 1886. Where the land-uplift is low, as around the coasts of southern Sweden, the sea level has risen by about 20 centimeters since 1886.



**Figure 6.** Sea levels corrected for the absolute land uplift (isostatic adjustment). Blue bars show the annual sea level averaged for 14 Swedish sea level records, compared to the 1886 level. The black and red line shows the gauss-filtered average and Stockholm sea level, respectively.

## International data exchange

Both real-time data and delayed mode data are routinely made available through several national and international programmes (Table 2). Real-time data have undergone gross error checking only, using a standard quality control protocol. Delayed mode has been screened and quality controlled using the procedures described by GLOSS, SeaDataNet and MyOcean. Real-time data can be obtained via: [http://www.smhi.se/en/Weather/Sweden-weather/Ocean-observations/havsvst\\_en.htm](http://www.smhi.se/en/Weather/Sweden-weather/Ocean-observations/havsvst_en.htm)

Programme	Data host	Frequency	Resolution	QC	Media	Notes
PSMSL	POL	Yearly	Month	Yes	Mail	All stations (23)
ESEAS	BODC	Monthly	HiRes*	Yes	FTP	All stations (23)
MYOCEAN	IFREMER	Daily	HiRes*	Yes	FTP	All stations (23)
GLOSS	VLIZ	Hourly	HiRes*	Yes	FTP	GLOSS stations (3)
BOOS/NOOS	SMHI	Hourly	Hour	No	FTP	All stations (23)
SEPRISE	SMHI	Hourly	Hour	No	FTP	All stations (23)
<a href="http://www.smhi.se">www.smhi.se</a>	SMHI	Hourly	Hour	No	www	Real-time stations (19)
<a href="http://www.boos.org">www.boos.org</a>	DMI	Hourly	Hour	No	www	Real-time stations (19)

\* 10-minute values and hourly maximum and minimum values. Minute-values are available for some periods, especially during storm periods.

**Table 2.** Sea level data are routinely made available through these programmes.

## The BOOS Data Exchange

The exchange of oceanographic data in the Baltic Sea is for the time being very well developed. Within the BOOS (Baltic Operational Oceanographic System) community we have developed an easy FTP-box system for exchange of data between the different institutions on a routinely basis (usually every hour). The resolution of the data is from about 5 minutes up to several hours, with the highest resolution for sea level data. SMHI is responsible for coordination of the data exchange and to implement routines for real-time quality control, validation and distribution of all sea level data coming from the Baltic Sea. For the moment, the system consists of 92 sea level stations (Figure 7). Together with other institutes in Europe we have also developed harmonized ways to exchange data in different EU-projects, such as MyOcean and SeaDataNet. This work will continue in the following years. A web page with station information and other metadata has been developed:

[http://www.smhi.se/hfa\\_coord/BOOS/dbkust/BOOS\\_Sealevel\\_Stations.html](http://www.smhi.se/hfa_coord/BOOS/dbkust/BOOS_Sealevel_Stations.html)



**Figure 7.** Sea level stations available through the BOOS Cooperation.

## New initiatives

During 2011 have SMHI installed new systems measuring sea level at three sites; Åhus, Ängelholm and Uddevalla. All three places indicates higher sea level during severe storm events, so it is very interesting to validate our model at these locations. Åhus is located in the Baltic, just north of Simrishamn (Figure 1). Ängelholm is located in the Laholm Bay in the Kattegat and Uddevalla is located north of Stenungsund in the Skagerack. At all places are we testing pressure sensors, where sea level is adjusted for salinity and water temperature variations (also recorded). In Uddevalla we are also testing a radar sensor. The stations are delivering high-resolution data every minute.



**Appendix 1.** Sea level stations owned and operated by the Swedish Maritime Administration. Real-time data from these stations can be obtained via:  
<http://vivakarta.sjofartsverket.se>

Station	Latitude	Longitude
Kalix	65° 47' 21" N	23° 18' 04" E
Larsgrund	65° 32' 56" N	22° 14' 20" E
Skellefteå	64° 40' 33" N	21° 17' 17" E
Holmsund	63° 41' 45" N	20° 20' 50" E
SkagsUdde	63° 11' 27" N	19° 00' 47" E
Svanö	62° 53' 24" N	17° 52' 09" E
Spikarna	62° 21' 48" N	17° 31' 51" E
Iggesund	61° 37' 28" N	17° 07' 44" E
Ljusne Orrskärskajen	61° 12' 25" N	17° 08' 44" E
Bönan	60° 44' 20" N	17° 19' 04" E
Loudden	59° 20' 33" N	18° 08' 29" E
Nynäshamn	58° 55' 02" N	17° 58' 07" E
Södertälje Saltsjön	59° 11' 31" N	17° 37' 56" E
Landsort	58° 44' 35" N	17° 52' 02" E
Vinterklasen (Oxelösund)	58° 39' 41" N	17° 07' 31" E
Juten	58° 38' 03" N	16° 19' 29" E
Västerbådan	57° 44' 50" N	16° 44' 31" E
Visby	57° 38' 22" N	18° 17' 06" E
Slite	57° 42' 21" N	18° 48' 37" E
Kalmar	56° 39' 32" N	16° 22' 42" E
Karlshamn	56° 09' 20" N	14° 49' 15" E
Flinten16	55° 33' 40" N	12° 48' 34" E
Flinten7	55° 35' 22" N	12° 50' 40" E
Malmö Hamn	55° 37' 07" N	12° 59' 06" E
Helsingborg	56° 02' 41" N	12° 41' 14" E
Halmstad	56° 38' 59" N	12° 50' 33" E
Varberg	57° 06' 34" N	12° 14' 30" E
Vinga	57° 37' 53" N	11° 36' 31" E
Måvholmsbådan	57° 40' 20" N	11° 42' 27" E
Karet	57° 41' 16" N	11° 52' 11" E
Marstrand	57° 53' 13" N	11° 35' 37" E
Brofjorden	58° 20' 10" N	11° 24' 17" E